



Distributed in the Interest  
of Product Development

# VANDERBILT *Report*

## Crop Protection Pesticide Formulations No. 921

**R.T. Vanderbilt Company, Inc.**  
**30 Winfield Street, P.O. Box 5150, Norwalk, CT 06856-5150**  
**Telephone: (203) 853-1400**  
**Fax: (203) 853-1452, Web Site: [www.rtvanderbilt.com](http://www.rtvanderbilt.com)**

Before using, read, understand and comply with the information and precautions in the Material Safety Data Sheets, label and other product literature. The information presented herein, while not guaranteed, was prepared by technical personnel and, to the best of our knowledge and belief, is true and accurate as of the date hereof. No warranty, representation or guarantee, express or implied, is made regarding accuracy, performance, stability, reliability or use. This information is not intended to be all-inclusive, because the manner and conditions of use, handling, storage and other factors may involve other or additional safety or performance considerations. The user is responsible for determining the suitability of any material for a specific purpose and for adopting such safety precautions as may be required. R.T. Vanderbilt Company, Inc. does not warrant the results to be obtained in using any material, and disclaims all liability with respect to the use, handling or further processing of any such material. No suggestion for use is intended as, and nothing herein shall be construed as, a recommendation to infringe any existing patent, trademark or copyright or to violate any federal, state or local law or regulation.

## Crop Protection Pesticide Formulations

The purpose of formulating pesticide active ingredients for crop protection is to uniformly spread a small amount of an active chemical over a large area. The goal is to ensure safety in handling and during application and to optimize pesticide efficacy. This requires that the pesticide formulation be chemically stable and physically uniform under all foreseeable storage conditions so that the minimum effective amount can be accurately applied to target areas.

R.T. Vanderbilt Company, Inc. supplies a variety of chemicals and minerals to assist the pesticide formulator. These include:

**Suspension Stabilizers:** **VAN GEL<sup>®</sup>** Magnesium Aluminum Silicate  
**VEEGUM<sup>®</sup>** Magnesium Aluminum Silicate

<b>VAN GEL B</b>	The standard economical grade for most suspensions
<b>VAN GEL ES</b>	The most electrolyte tolerant grade
<b>VEEGUM</b>	The standard grade for a wide range of applications
<b>VEEGUM CER</b>	Smectite clay/CMC blend; high efficiency suspension stabilizer
Several additional grades are available to match formulation requirements.	

**VANZAN<sup>®</sup>** Xanthan Gum

<b>VANZAN</b>	The general purpose grade suitable for most applications
<b>VANZAN D</b>	Surface-treated to facilitate dispersion without lumping, then rapid dissolution at pH >9

**Dispersants:** **DARVAN<sup>®</sup>** Dispersant

<b>DARVAN 2</b>	Sodium lignosulfonate
<b>DARVAN 670</b>	Sodium salts of polymerized alkyl naphthalene sulfonic acid
Several additional grades are available to match formulation requirements.	

**Diluents:** **CONTINENTAL<sup>®</sup>** Clay  
**PYRAX<sup>®</sup>** Pyrophyllite

<b>CONTINENTAL</b>	Agricultural grade kaolin with unusually fine particle size
<b>PYRAX ABB</b>	An inert diluent for pesticide powders

## LIQUID CONCENTRATES

**Aqueous flowables (AF)** are concentrated 40% to 70% w/w suspensions of micronized insoluble active pesticide in water. Prior to spraying on target areas, aqueous flowables are diluted with water in a spray tank to achieve the minimum effective pesticide concentration. AFs must be formulated for low viscosity and good fluidity, so that transfer to the spray tank is easy and complete. This requires an effective wetting agent and an efficient dispersing agent to ensure adequate dispersion of the pesticide in the water. Since the active ingredients in AFs are insoluble, good suspension stability is essential. If the suspension settles and leaves sediment at the bottom of the container, the application of the pesticide may be too weak to be effective. Further, disposal of the residue in the container becomes a problem. A combination of smectite clay (aka bentonite) and xanthan gum works synergistically to provide excellent long term suspension stability at low viscosity and at low cost.

There are three techniques commonly used to stabilize aqueous flowables:

1. When the pesticide is already milled to the desired particle size, the smectite clay and the organic gum (xanthan gum, cellulose gum), if used, are first thoroughly hydrated before the other ingredients are added with a dispersing mixer.
2. When the pesticide needs to be milled, a pre-suspension of the smectite clay can be formed as above, reserving the organic gum. This is then media milled (attritor, ball mill, sand mill) until the pesticide particles are reduced to the desired size. The organic gum is added at the end of the milling cycle, avoiding shear degradation.
3. All ingredients except the organic gum are added directly to the media mill without pre-dispersion. The smectite clay hydrates as the pesticide particles are milled. Generally, smectite clay is best delaminated when it is pre-hydrated, but the high shear imparted by the mill is usually sufficient. The organic gum, typically xanthan gum, is added and dissolved as the last step; it interacts with the clay to form the colloidal structure that keeps the pesticide particles uniformly dispersed and suspended during storage.

The aqueous flowable formulas in Table 1 demonstrate the use of **DARVAN** dispersants with the **VAN GEL B / VANZAN** suspending agent combination to produce low viscosity fluid concentrates with excellent physical stability. These formulas were prepared in 2 kg batches in a laboratory impact mill. All ingredients were charged to the mill except the xanthan gum. Milling proceeded until the average particle size of the active was reduced to less than 5 micrometers. The xanthan gum was then added and milling was continued only long enough to allow the gum to dissolve.

**Emulsifiable Concentrates (EC)** are blends of pesticide, emulsifiers and adjuvants dissolved in a volatile oil. Low melting point or liquid pesticide actives have traditionally been formulated into EC's. When the EC is added to water in the spray tank, it forms a stable dilute emulsion. ECs are a convenient means of using water as a vehicle for oil-soluble pesticides. However, their oily solvent base creates several hazards: ECs can be absorbed through the skin, burn foliage, and attack the rubber and plastic parts of spray equipment.

**Concentrated emulsions (EW)** contain up to 50% w/w of oil-soluble pesticide. EWs minimize the disadvantages of emulsifiable concentrates by minimizing the level of solvent needed to dissolve the pesticide, and then emulsifying this solution in water. The concentrated emulsion is simply diluted in the spray tank for application. The combination of smectite clay and xanthan gum stabilizes the concentrated emulsion against separation in the same way that it stabilizes concentrated AF suspensions. To prepare the concentrated emulsion, the smectite clay is hydrated before the emulsion is formed. The xanthan gum can be dissolved in the water either before or after emulsification.

**Suspo-emulsions (SE)** are produced by combining an aqueous flowable with a concentrated emulsion. The need to ensure long-term fluidity and physical stability requires that particular care be taken in the selection of proper wetting agents and emulsifiers. The smectite clay/xanthan gum (e.g., **VAN GEL B/VANZAN**) combination ensures the stability of the dispersed oil phase against coalescence and prevents the agglomeration of pesticide particles. **DARVAN** dispersants are useful in promoting the optimum dispersion of the pesticide particles.

**Table 1. Aqueous Flowable Formulas**

	<b>No. 388 Carbaryl 4 lbs/gal</b>	<b>No. 386 Carbaryl 5 lbs/gal</b>	<b>No. 387 Atrazine 4 lbs/gal</b>	<b>No. 389 Sulfur 6 lbs/gal</b>
<b>VAN GEL® B</b> Magnesium Aluminum Silicate	0.40%	0.40%	0.40%	0.50%
<b>DARVAN® 2</b> Dispersant	2.00	3.50	—	—
<b>DARVAN 670</b> Dispersant	—	—	2.00	2.50
Water	41.24	25.78	41.42	34.09
Propylene Glycol	10.00	15.00	7.50	9.50
Surfynol® TGE <sup>1</sup>	0.10	0.10	—	0.10
Surfynol 104H <sup>1</sup>	—	—	0.50	—
Atlox® 4896 <sup>2</sup>	—	—	2.00	—
Triton® X114 <sup>3</sup>	—	—	—	0.15
Citric acid	0.08	0.08	—	—
Triethanolamine	—	—	qs, pH 7	—
Preservative	0.10	0.10	0.10	0.10
Carbaryl, 99% Tech.	46.00	55.00	—	—
Atrazine, 98% Tech.	—	—	46.00	—
Flour Sulfur	—	—	—	53.00
<b>VANZAN®</b> Xanthan Gum	0.08	0.04	0.08	0.06
<b>Viscosity, cps<sup>A</sup></b>				
1 Day	480	900	420	800
1 Month	460	880	420	780
3 Months	460	870	410	760
<b>Stability, 3 months</b>				
Gelation, 4/20/50°C	none	none	none	none
Bleed, 4/20/50°C	0/tr/1%	tr/tr/1%	0/0/tr	0/3%/tr
Packing, 4/20/50°C	none	none	none	none
<b>Freeze/Thaw, 5 cycle</b>	passed	passed	passed	passed
<b>Bloom</b>	passed	passed	passed	passed
<b>Average particle size, µm<sup>B</sup></b>	3.1	4.5	2.7	2.7
<b>pH</b>	5.5	5.5	7.0	7.3
<b>Density, lbs/gal</b>	9.2	9.4	9.1	11.6

<sup>A</sup>Brookfield, 60 rpm <sup>B</sup>Coulter

<sup>1</sup>Air Products and Chemicals, Inc., Allentown, PA

<sup>2</sup>ICI Americas, Inc., Bridgewater, NJ

<sup>3</sup>Union Carbide Chemicals & Plastics Technology Corporation, Danbury, CT

## WETTABLE POWDERS

**Wettable powders (WP)** are dry concentrates containing 50% or more of micronized active pesticide blended with a wetting agent, a dispersing agent and a finely ground diluent. Prior to spraying on target areas, wettable powders are dispersed into water in a spray tank to achieve the minimum effective pesticide concentration. The active may be a crystalline solid or a liquid or low melting point compound absorbed onto a carrier. The wetting agent ensures easy dispersion of the active in water, and the dispersing agent, such as a **DARVAN** dispersant, inhibits the agglomeration of pesticide particles. **CONTINENTAL** clay is widely used as the solid diluent in wettable powders because of its naturally fine particle size, its broad compatibility with pesticide actives and its ability to inhibit the settling of pesticide particles in the spray tank. Typical formulations are shown in Table 2.

**Table 2. Typical Wettable Powder Formulations**

Atrazine	50%	—	—	—	—
Carbaryl	—	75%	—	—	—
Diuron	—	—	80%	—	—
Mancozeb	—	—	—	75%	—
Sulfur	—	—	—	—	80%
Morwet <sup>®</sup> EFW <sup>1</sup>	4-6	4-6	4-6	4-6	4-6
<b>DARVAN<sup>®</sup> 2</b> Dispersant	—	4-6	—	—	—
<b>DARVAN 670</b> Dispersant	3-4	—	4-6	4-6	4-6
<b>CONTINENTAL<sup>®</sup></b> Clay	qs to 100	qs to 100	qs to 100	qs to 100	qs to 100

<sup>1</sup>Akzo Nobel Surface Chemistry LLC, Chicago, IL

Because wettable powders are applied as dilute aqueous suspensions, drift into non-target areas is more easily controlled than with dusts, but exposure to respirable pesticide powder remains a concern.

## WATER DISPERSIBLE GRANULES

**Water dispersible granules (WDG), or dry flowables (DF),** are wettable powders that have been aggregated into uniform granules for easier handling and to eliminate respirable particles. They are dispersed in the spray tank and applied as dilute suspensions in the same way as wettable powders. WDGs incorporate the same ingredients as wettable powders, including **DARVAN** dispersants and **CONTINENTAL** clay, although generally with a higher level of active and less diluent. The granules must be strong enough to resist crumbling into powder, yet still readily and completely disintegrate and disperse in the spray tank. This avoids nozzle clogging and ensures the finest particle size of the active ingredient for optimum efficacy.

Several methods are used to form granules from the starting powder blend. These include pan granulation, fluid bed granulation, spray drying, high speed mixer agglomeration and extrusion granulation. Extrusion granulation is generally preferred on the basis of safety, versatility and economy.

## DUSTS

**Dusts (D)** are formulated as ready-to-use pesticides, with no dilution required. The active is either a crystalline solid ground to 1-10 micrometers, or a liquid or waxy compound absorbed onto an inert mineral carrier. The concentration of active is typically less than 10% by weight, with a finely ground mineral diluent making up the balance. Aerial application of dusts was once commonly practiced, but this has all but been abandoned due to inhalation hazards and the tendency of dusts to drift into non-target areas. Today dusts are used for small area treatment such as gardens, for seed treatments and for the control of parasites on pets and livestock.

The preferred diluent for pesticide dusts is **PYRAX ABB** pyrophyllite. This finely ground pyrophyllite is an inert, non-hygroscopic, neutral pH mineral which provides optimal compatibility with pesticide actives, and is safe for plants and animals. **PYRAX ABB** has a relatively high bulk density, resulting in “heavy” dusts that drift less on application to plants, and that enable animal dusts to penetrate fur for good skin contact.

rev.02/08

**VAN GEL, VEEGUM, DARVAN, VANZAN, PYRAX** and **CONTINENTAL** are registered trademarks of R.T. Vanderbilt Company, Inc.

Atlox is a registered trademark of ICI Americas, Inc.

Surfynol is a registered trademark of Air Products and Chemicals, Inc.

Morwet is a registered trademark of Akzo Nobel Surface Chemistry LLC

Triton is a registered trademark of Union Carbide Chemicals & Plastics Technology Corporation

**For additional information regarding our  
high quality minerals and chemicals,  
please visit our website:**

**[www.rtvanderbilt.com](http://www.rtvanderbilt.com)**

- Technical data sheets
- MSDS information
- Sample requests
- Specifications
- Product brochures
- Articles
- Presentations
- Reports